

Flavonoid and Flavonoid glycoside from *Butea superba* Roxb. and their cAMP Phosphodiesterase Inhibitory Activity

Sophon Roengsumran,¹ Amorn Petsom,¹ Nattaya Ngamrojanavanich,¹ Thanatip Rugsilp,¹ Pailin Sittiwicheanwong,¹ Prapas Khorphueng,¹ Wichai Cherdshewasart,² and Chaiyo Chaichantipyuth³

A flavonoid (3, 7, 3'-Trihydroxy-4'-methoxyflavone) (**1**) and a flavonoid glycoside (3, 3'-dihydroxy-4'-methoxyflavone-7-O- β -D-glucopyranoside) (**2**) were isolated from the tuber root of *Butea superba* Roxb. The structures were determined on the basis of spectral analysis, including 2D-NMR techniques. These compounds show higher inhibitory effects on cAMP phosphodiesterase than caffeine and theophylline.

Key Words: *Butea superba*, flavonoid, flavonoid glycoside, cAMP phosphodiesterase inhibition

¹ Department of Chemistry, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand.

² Department of Biology, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand.

³ Department of Pharmacognocoy, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand.

ฟลาวอนอยด์ และฟลาวอนอยด์ไกลโคไซด์จากกวาวเครือแดง และ ฤทธิ์ต่อต้านไซคลิกเอเอ็มพีฟอสโฟไดเอสเทอร์

โสภณ เริงสำราญ, อมร เพชรสม, นาทยา งามโรจนวนิชย์, ธนาธิป รักศิลป์,
ไพลิน สิทธิวิเชียรวงศ์, ประภาส ขอฟุ้ง, วิชัย เชิดชูวิศาสตร์ และ ชัยโย ชัยชาญทิพยุทธ (2543)
วารสารวิจัยวิทยาศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย 25(1)

ฟลาวอนอยด์ (3, 7, 3'-ไฮดรอกซี-4'-เมธอกซีฟลาโวน) (1) และฟลาวอนอยด์
ไกลโคไซด์ (3, 3'-ไดไฮดรอกซี-4'-เมธอกซีฟลาโวน-7-O- β -D-กลูโคไพราโนไซด์) (2) แยกได้
จากส่วนรากของกวาวเครือแดง ได้ทำการหาสูตรโครงสร้างของสารดังกล่าวโดยอาศัยผลการ
วิเคราะห์ทางสเปกโทรสโกปี ซึ่งประกอบด้วยเทคนิค 2D-NMR สารประกอบทั้งสองชนิด
แสดงผลในการยับยั้งไซคลิกเอเอ็มพี ฟอสโฟไดเอสเทอร์อย่างแรง

คำสำคัญ กวาวเครือแดง ฟลาวอนอยด์ ฟลาวอนอยด์ไกลโคไซด์ การยับยั้ง
ไซคลิกเอเอ็มพีฟอสโฟไดเอสเทอร์

INTRODUCTION

Butea superba Roxb. is a plant in the Family Papilionaceae and has the characteristics of being a climber that wraps itself around large trees. One branch has 3 leaves, the flowers are of a yellowish orange color and the plant grows out in the open. The long tuber root of the plant is buried under the ground like the tuber root of a yam. This plant reproduces through seeds and the propagation of its tuber root. This plant can be found growing in forests in Thailand's northern regions, eastern regions and along Kanchanaburi Province. The tuber and stem of the plant are used in medicines believed to

give strength and power and increase male sexual performance. Thus, this plant has come to be known as one type of miracle herb. Since *Butea superba* Roxb. helps to enhance human health, it is therefore very interesting to investigate the chemical constituents of this plant and their biological activity. The bioactivity of each constituent was tested for an inhibitory effect towards cAMP phosphodiesterase, which has been shown to be important in controlling bodily function and involved a wide number of diseases [1].

Table 1. ¹H NMR spectral data of compounds 1 and 2 (500 MHz DMSO).

Position	1 δH (J in Hz)	2 δH (J in Hz)
2		
3		
4		
5	7.96 d (8.8)	8.05 d (8.8)
6	6.94 dd (2.1, 8.8)	7.15 dd (2.1, 8.8)
7		
8	6.85 d (2.1)	7.22 d (2.4)
9		
10		
1'		
2'	8.35 d (2.1)	8.40 d (2.0)
3'		
4'		
5'	6.98 d (8.8)	6.98 d (8.8)
6'	7.50 dd (2.1, 8.8)	7.52 dd (2.1, 8.8)
-OCH ₃	3.79 s	3.78 s
1"		5.10 d (7.6)
2"		3.30 dd (7.6, 9.5)
3"		3.35 t (9.5)
4"		3.18 t (9.5)
5"		3.45 m
6"a		3.45 dd (12.0, 2.5)
6"b		3.70 dd (12.0, 4.5)

Table 2. ^{13}C NMR (125 MHz DMSO) and 2D Long-Range ^1H - ^{13}C Correlations in the HMBC Spectra of Compounds **1** and **2**.

Position	1		2	
	δC	HMBC	δC	HMBC
2	158.9		159.1	
3	146.8		147.0	
4	174.6		174.8	
5	127.3	4, 7, 9	127.1	4, 7, 9
6	115.2	5, 7, 8, 10	115.7	5, 7, 8, 10
7	162.6		161.5	
8	102.1	6, 7, 9, 10	103.5	6, 7, 9, 10
9	157.4		157.1	
10	116.6		118.5	
1'	124.2		123.5	
2'	153.4	1', 3', 4', 6'	153.6	1', 3', 4', 6'
3'	157.3		157.1	
4'	146.9		146.6	
5'	113.6	1', 3', 4'	113.7	1', 4', 6'
6'	130.0	1', 2, 4', 5'	130.1	1', 3', 4', 5'
-OCH ₃	55.2		55.2	
1''			100.1	
2''			73.2	
3''			76.5	
4''			69.8	
5''			77.2	
6''			60.8	

EXPERIMENTAL

General Experimental Procedures

All commercial grade solvents were distilled prior to use. Melting points were determined on a Fisher-Johns melting point apparatus and are reported uncorrected. The optical rotation was determined on a JASCO DIP-370 digital polarimeter. Measurements of UV spectra were carried out on a Milton-Roy Spectronic 3000 Array UV/VIS spectrophotometer. IR spectra were recorded on a Perkin-Elmer model 1760X FT-IR spectrophotometer. Spectra of solid samples were recorded as KBr pellets. The ^1H and ^{13}C NMR spectra were recorded at 500.00 and 125.65 MHz, respectively, on a JEOL JNM-A500 NMR spectrometer. LREIMS

were obtained with a Fisons Instruments model Trio 2000 mass spectrometer at 70 eV.

Plant Materials

The tubers of *Butea superba* Roxb. were collected from Amphur Muang, Lumpang Province, Thailand in May 1997. Botanical identification was claimed through comparison with a voucher specimen No. BKF 70163 in the herbarium collection of Royal Forest Department of Thailand.


Extraction and Isolation

Powdered sun dried roots (16.0 kg) of *Butea superba* were repeatedly extracted with MeOH (5x10 L). The MeOH extracts were filtered and evaporated under reduced pressure to obtain a dark-red gummy residue

(106.0 g) of MeOH crude extract. This MeOH crude extract was subsequently re-extracted with hexane and then CHCl_3 to leave the final insoluble residue (72.0 g). The hexane and CHCl_3 extract fractions were evaporated under reduce pressure to give a hexane crude extract (21.0 g) and CHCl_3 crude extract (12.0 g), respectively. The CHCl_3 crude extract (12.0 g) was subjected to silica gel column chromatography using gradient elution with CHCl_3 and MeOH in a stepwise fashion. Compound **1** was eluted with 5% MeOH in CHCl_3 . Similar fractions were combined and the solvent was removed under reduced pressure to give compound **1** (135.0 mg) after recrystallization from MeOH and CHCl_3 . The final residue (72.0 g) was separated by column chromatography on Silica gel using gradient elution with increasing amounts of MeOH in CHCl_3 to give compound **2** (60.0 mg) from 10% of MeOH in CHCl_3 fraction. Compound **2** was recrystallized from MeOH and CHCl_3 .


Flavonoid (3, 7, 3'-Trihydroxy-4'-methoxyflavone) (compound **1**): pale yellow needle crystal, mp 258-260°C (d) [lit 288(d), but no spectroscopic data for verification], (found: C, 64.0; H, 3.9, $\text{C}_{16}\text{H}_{12}\text{O}_6$ required: C, 64.0; H, 4.0); UV λ_{max} EtOH nm 254, 316 and 365; IR $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} ; 3340-3000, 2940, 1650, 1594, 1575, 1500, 1450, 1380, 1260, 1090, 1020, 790; ^1H and ^{13}C -NMR Table 1; EIMS m/z (rel. int.) 300[M^+] (25), 282 (30), 268 (100), 253 (25), 132 (60).

Flavonoid glycoside (3,5'-Dihydroxy-4'-methoxyflavone-7-O- β -D-glucopyranoside) (compound **2**); white amorphous solid; mp 210-212°C, (found: C, 57.0; H, 4.5, $\text{C}_{22}\text{H}_{22}\text{O}_{11}$ requires: C, 57.1; H, 4.8); $[\alpha]_{\text{D}}^{25} +9.5$ (c 1.05, MeOH); UV λ_{max} EtOH nm 267, 290 and 355; $\nu_{\text{max}}^{\text{KBr}}$ cm^{-1} ; 3600-3100, 2900, 1650, 1550, 1450, 1300, 1260, 1060-1030, 891, 800; ^1H and ^{13}C -NMR Table 1 and 2; EIMS m/z (rel. int.) 282 (20), 268 (100), 253 (23), 133 (12), 132 (90).



1 R = H

Scheme 1



Scheme 2

Bioassay

Phosphodiesterase activity was determined from the amount of inorganic phosphate liberated during the reaction by the malachite green method [2, 3]. Phosphodiesterase assay solutions were prepared as follows: (a) the enzyme solution contains phosphodiesterase (0.037 unit mL^{-1}), 5-nucleotidase (1.67 unit mL^{-1}), MgCl_2 (5 mM) and Tris HCl (0.2 M); (b) reaction mixture A contains malachite green (0.33 mM), polyvinyl alcohol (3.87g L^{-1}) and ammonium molybdate in 6 M HCl (8.33 mM). The test sample was dissolved in 1.5 % dimethyl sulfoxide in water. The reaction was performed by the addition of cAMP (10 mM, 100 μL) to the enzyme solution (400 μL) at 30°C. Then the sample solution (500 μL), reagent mixture A (1.0 mL) and 25% sodium citrate (200 μL) were added to the above solution successively at 5 minute intervals. The absorbance of the colour complex was

observed at 630 nm using a UV/VIS spectrophotometer referred against a mixed reagent blank. A calibration curve, obtained by this procedure using potassium dihydrogen phosphate solution of a known concentration, was used to determine the amount of phosphate present in the assay. For control experiment, dimethyl sulfoxide was added instead of the solution of the sample to minimize the effects of the solvent and theophylline and caffeine were used as positive controls for phosphodiesterase assay. The IC₅₀ values of compound (1) and (2) were determined from the calibration curve of sample concentration against cAMP phosphodiesterase activity.

RESULTS AND DISCUSSION

The 3, 7, 3'-Trihydroxy-4'-methoxyflavone (compound 1) was obtained from a chloroform soluble crude extract from the root of *Butea superba* by silica-gel column chromatography using a gradient elution with chloroform and methanol. The IR spectrum of 1 showed a broad absorption band between 3000 and 3340 cm⁻¹ of OH stretching and a strong absorption band at 1650 cm⁻¹, which was consistent with a conjugated carbonyl group. The carbon-carbon double bond stretching vibration of the aromatic phenyl group was also observed at 1594, 1575 and 1500 cm⁻¹. The UV spectrum exhibited absorption maxima at 254 and 365 nm which are characteristic absorption bands of a flavone skeleton [4]. The molecular formula of C₁₆H₁₂O₆ was established for compound 1 from the elemental analysis, LREIMS and ¹H and ¹³C NMR data (Tables 1 and 2). The ¹H and ¹³C NMR spectra together with 2D NMR experiments allowed the complete structure of compound 1 to be established. The occurrence of flavonoid (1) was clearly determined from the ¹H 500 MHz NMR spectrum which displayed six aromatic protons at δ 6.85 (d, J=2.1 Hz), 6.94 (dd, J=2.1, 8.8 Hz), 6.98 (d, J=8.8 Hz), 7.50 (dd, J=2.1, 8.8 Hz), 7.96 (d, J=8.8 Hz), and 8.35 (d, J=2.1 Hz) and one methyl proton singlet

at 3.79 ppm. Detailed analysis of the 2D ¹H and ¹³C NMR spectrum, including COSY, NOESY, HMQC and HMBC supported the structure of 3, 7, 3'-Trihydroxy-4'-methoxyflavone (1). Although compound 1 is a known compound isolated as a minor constituents of quebracho tannin extract [5], its ¹H and ¹³C NMR spectral data have not been published before.

The 3, 3'-Dihydroxy-4'-methoxyflavone-7-O-β-D-glucopyranoside (compound 2) was isolated as an amorphous white solid powder from MeOH extract residue by silica gel column chromatography using CHCl₃-MeOH as the eluent. The IR spectrum of compound 2 displayed absorption at 3100-3600 cm⁻¹ of OH stretching, and strong absorption at 2900 cm⁻¹ of C-H stretching. The absorption band of a conjugated carbonyl group appeared at 1650 cm⁻¹. The molecular formula of compound 2 was assigned as C₂₂H₂₂O₁₁ based on elemental analysis and ¹H and ¹³C NMR (Table 1 and 2) while its EI-MS showed no molecular ion peak pointing to its glycoside nature. The ¹H NMR spectrum of compound 2 in DMSO showed the presence of a sugar moiety by one proton doublet at δ 5.10 (J=7.6 Hz, H-1''), one proton doublet of doublets at δ 3.30 (J=7.6, 9.5 Hz, H-2''), one proton triplet at δ 3.35 (J=9.5 Hz, H-3''), one proton triplet at δ 3.18 (J=9.5 Hz, H-4'') and one proton multiplet at δ 3.45 (H-5''). Two protons of C-6'' included one proton showed doublet of doublets at δ 3.45 (J=12.0, 2.5 Hz, H-6''a) and another one proton showed doublet of doublets at δ 3.70 (J=12.0, 4.5 Hz, H-6''b). The chemical shifts at δ 6.98, 7.15, 7.22, 7.52, 8.05 and 8.40 were assigned to flavonoid protons at C-5', C-6, C-8, C-6', C-5 and C-2' respectively, on the basis of their similarity to signals observed for compound 1.

Their ¹H and ¹³C-NMR spectra indicate that glycoside 2 has a glycone portion identical to that in compound 1. In the ¹H NMR spectra (Table 1), an unusual pattern of 7-O-glycosylation was indicated by downfield shifts of H-6 (ca. +0.21 ppm) and H-8 (ca. +0.36 ppm) with respect to

compound 1. Similarly, in the ^{13}C -NMR spectra of compound 2 (Table 2), 7-O-glycosylation was confirmed by the diagnostic [6] upfield shift of C-7 (-0.24 ppm) and by downfield shifts of the ortho-related C-8 (+1.45 ppm) and C-6 (+0.55 ppm) and para-related C-10 (+1.99 ppm) carbon with respect to compound 1. The ^1H and ^{13}C NMR data indicated the β -configuration at an anomeric position for the glucopyranosyl unit (Table 1 and 2). Therefore, the structure of compound 2 was assigned as 3,3'-dihydroxy-4'-methoxyflavone-7-O- β -D-glucopyranoside.

From the bioactivity testing of compound 1 and compound 2, it was found that both of these compounds were effective in inhibiting cAMP phosphodiesterase. These two compounds have $\text{IC}_{50} = 190$ and $58 \mu\text{g/mL}$, respectively. The cAMP phosphodiesterase inhibition of both these compounds was more effective than those of theophylline ($\text{IC}_{50} = 615 \mu\text{g/mL}$) and caffeine ($\text{IC}_{50} = 420 \mu\text{g/mL}$). The cAMP phosphodiesterase enzyme has a main function in the hydrolysis of the intracellular cAMP. Substances that inhibit cAMP phosphodiesterase are therefore capable of stimulating the functioning of the central nervous system (CNS) and stimulating the functioning of cells [7]. Thus, papaverin, dipyrindamole, caffeine and theophylline are effective when the cAMP phosphodiesterase is inhibited [8, 9]. Furthermore, substances that have an effect in inhibiting cAMP phosphodiesterase also take part in controlling numerous severe diseases, including diabetes [10-13], hypertension [14, 15], asthma [16], hepatomas [17], psoriasis [18] and possibly cancer [19, 20]. In addition, substances that inhibit phosphodiesterase are shown to have effects in controlling platelet-aggregation inhibition [21- 26].

Therefore, *Butea superba* Roxb. is composed of flavonoids that are effective in inhibiting the cAMP phosphodiesterase enzyme, which are very beneficial to the human body. At least, when taking this herb, the body will begin to feel healthier. This

herb will also help control the numerous diseases mentioned beforehand.

ACKNOWLEDGEMENTS

We thank the Graduate School and Department of Chemistry, Faculty of Science, Chulalongkorn University, for financial support and the staff of the Scientific and Technology Research Equipment Centre, Chulalongkorn University, for recording highfield NMR spectra and performing 2D NMR experiments. We also thank Mr. Thanase U. Patriyakul of Eiwlee Industrial for providing some plant materials.

REFERENCES

1. M. S. Amer, and W. E. Kreighbaum, *J. Pharmac. Sci.*, **64**, 1 (1975)
2. N. Chairungsrilerd, K. Takeuchi, Y. Ohizumi, S. Nozoe and T. Ohta, *Phytochemistry*, **43**, 1099 (1996)
3. K. M. Chan, D. Delfert and K. D. Janger, *Analyt. Biochem.*, **157**, 375 (1986)
4. K. R. Markham, *Techniques of Flavonoid Identification*, Academic Press; London, 36 (1982)
5. K. S. Kirby and T. White, *Biochem. J.*, **60**, 582 (1955)
6. M. D' Agostino, V. De Feo, F. De Simone, C. Pizza, *Phytochemistry*, **22**, 2440 (1991)
7. C. A. Smith and E. J. Wood, *Molecular and Cell Biochemistry: Cell Biology*, Chapman and Hall, 209 (1992)
8. I. Weinryb, M. Chasin, C. A. Free, D. N. Harris, H. Goldenberg, I. M. Michel, V. S. Paik, M. Phillips, S. Samaniego and S. M. Hess, *J. Pharmac. Sci.*, **61**, 1556 (1972)
9. T. Nikaido, T. Ohmoto, H. Noguchi, T. Kinoshita, H. Saitoh and U. Sankawa, *Phanta. Med.* **43**, 18 (1981)
10. J. G. Hemington, M. Chenoweth and A. Dunn, *Biochim. Biophys. Acta.*, **304**, 552 (1973)
11. I. Das and E. B. Chain, *Biochem. J.*, **128**, 95 (1972)

12. A. H. Kissebah and T. R. Fraser, *Horm. Metab. Res.*, **4**, 72 (1972)
13. S. J. Pilkis, H. J. Exton, R. A. Johnson, and C. R. Park, *Fed. Proc.*, **32**, 299 (1973)
14. M. S. Amer, *Science* **179**, 807 (1973)
15. L. Triner, Y. Vulliemoz, M. Verosky, W. M. Manger and G. G. Nahas, *Fed. Proc.*, **31**, 556 (1972)
16. C. W. Parker and J. W. Smith, *J. Clin. Invest.*, **52**, 48 (1973)
17. A. R. Rhoads, H. P. Morris and W. L. West, *Cancer Res.*, **32**, 2651 (1972)
18. R. K. Wright, S. H. Mandy, K. M. Halprin and S. L. Hsia, *Arch. Dermatol.*, **107**, 47 (1973)
19. P. Emmelot and C. J. Bos, *Biochim. Biophys. Acta*, **249**, 285 (1971)
20. V. Tomasi, A. Rethy and A. Trevisani, *Life Sci.*, **12**, 145 (1973)
21. M. Horlington and P. A. Watson, *Biochem. Pharmacol.*, **19**, 955 (1970)
22. R. L. Vigdahl, N. R. Marquis and P. A. Tavormina, *Biochem. Biophys. Res. Commun.*, **37**, 409 (1969)
23. R. R. Gorman, S. Bunting and O. V. Miller, *Prostaglandins*, **13**, 377 (1977)
24. S. M. Wolfe and N. R. Shulman, *Biochem. Biophys. Res. Commun.*, **35**, 265 (1969)
25. N. A. Meanwell, B. C. Pearce, H. R. Roth, E. C. R. Smith, D. L. Wedding, J. J. K. Wright, J. O. Buchanan, U. M. Baryla, M. Gamberdella, E. Gillespie, D. C. Hayes, S. M. Seiler, H. C. Stanton, G. B. Zavoico and J. S. Fleming, *J. Med. Chem.*, **35**, 2672 (1992)
26. N. A. Meanwell, P. Hewawasam, J. A. Thomas, J. J. K. Wright, J. W. Russell, M. Gamberdella, H. J. Goldenberg, S. M. Seiler and G. B. Zavoico, *J. Med. Chem.*, **36**, 3251 (1993)

Received: August 6, 1999

Accepted: February 1, 2000

